

BOTULIN

EVIL STUFF

PLEASE DO NOT MAKE THIS SUBSTANCE, IT IS EXTREMELY HAZARDOUS!!!

BOTULISM IS THE DISEASE CAUSED BY EATING FOOD IN WHICH THE BACTERIA CLOSTRIDIA BOTULINUM HAS GROWN.

THIS BACTERIA MAKES A POISONOUS PROTEIN CALLED BOTULIN.

BOTULIN IS CREATED AS A WASTE PRODUCT OF THE BACTERIA WHILE IT IS EATING THE FOOD. HYDROGEN SULFIDE GAS IS ALSO GIVEN OFF, WHICH SMELLS LIKE ROTTEN EGGS.

BOTULIN IS THE SECOND MOST POWERFUL POISON KNOWN. THE FATAL DOSE OF PURE BOTULIN POISON IS AROUND 1 MICROGRAM. THERE ARE 1 MILLION FATAL DOSES IN A GRAM OF PURE BOTULIN!!!

THE BACTERIA IS FOUND ALL OVER THE WORLD, IN ALMOST EVERY FEW INCHES OF SOIL, RICH IN ORGANIC MATERIALS. JUST GO OUT TO A MOIST SHADED, ORGANIC SOIL BED AND DIG A COUPLE INCHES DOWN AND YOU HAVE PROBABLY FOUND CLOSTRIDIA BOTULINUM BACTERIA.

THIS BACTERIA COMES IN AROUND 7 DIFFERENT STRAINS, ONLY A & E ARE FATAL TO HUMANS.

THIS BACTERIA IS ALSO ANAEROBIC, WHICH MEANS IT WILL ONLY GROW IN THE ABSENCE OF AIR.

BOTULISM SYMPTOMS:

WHEN INGESTED, SYMPTOMS OCCUR WITHIN 12-36 HOURS AND INCLUDE FATIGUE, DIZZINESS, HEADACHE, CONSTIPATION, VERTIGO, DIFFICULTY IN SWALLOWING AND IN SPEECH, REGURGITATION OF FLUIDS FROM THE NOSE AND MOUTH, MUSCULAR INCOORDINATION, RELEASE OF BLADDER, EVENTUALLY DEATH FROM RESPIRATORY FAILURE.

WHEN INJECTED OR TAKEN THROUGH THE BLOOD STREAM, DEATH IS QUICK AND SYMPTOMLESS.

BOTULIN TOXIN CAN BE MIXED WITH DMSO AND SPRAYED ON DOORKNOBS AND WILL ABSORB INTO THE SKIN AND CAUSE QUICK DEATH.

THE BACTERIA THAT MAKES BOTULIN IS FOUND ALL OVER THE WORLD. A RANDOMLY CHOSEN SOIL SAMPLE IS LIKELY TO CONTAIN QUITE A FEW SPORES OF THIS BACTERIA. SPORES ARE LIKE SEEDS FOR BACTERIA, AND CAN WITHSTAND VERY HARSH TREATMENT. THIS PROPERTY WILL COME IN VERY HANDY IN ANY ATTEMPT TO GROW BOTULISM GERMS, BECAUSE OTHER GERMS CAN BE WIPED OUT BY HEATING IN HOT WATER, LEAVING THE SPORES TO GERMINATE AND TAKE OVER ONCE THEY COOL DOWN.

BOTULISM GERMS CAN NOT SURVIVE EXPOSURE TO AIR. OXYGEN KILLS THEM BUT DOES NOT KILL THEIR SPORES. IT ALSO DOES NOT KILL THE POISON WHICH HAS ALREADY BEEN PRODUCED, HEAT WILL KILL THE POISON.

IDEAL CONDITIONS FOR CULTIVATING THE BACTERIA:

THEY INCLUDE A FOOD SOURCE RICH IN PROTEIN, A pH THAT IS NEARLY NEUTRAL (IDEALLY AROUND 7.2), A TEMPERATURE AT 90 DEG F OR 35 DEG C. NO WARMER OR YOU MAY DAMAGE THE BACTERIA.

WHERE DO YOU FIND THIS BACTERIA???

THE BACTERIA PREFERS DARK, RICH, FAIRLY WET SOIL THAT IS RICH IN ORGANIC MATTER. IT IS MORE POPULATED IN WARM AREAS, TYPE A BACT IS THE PREDOMINANT TYPE OF GERM IN THE US. VIRGIN, UNTURNED SOIL HOLDS THE GREATEST PREPONDERANCE OF TYPE A GERMS. IT IS HARDER TO FIND SOIL SAMPLES THAT DO NOT CONTAIN THE GERM. GO OUT AND COLLECT SEVERAL SAMPLES OF SOIL FROM DIFFERENT LOCATIONS, MAKE SURE THE SAMPLES COME FROM 3-4 INCHES BELOW THE SURFACE.

LETS ISOLATE SOME GOOD STRAINS...

TAKE ABOUT 3 SAMPLES OF ABOUT 1 TABLESPOON EACH, TRY TO KEEP THEM IN THEIR ORIGINAL CONDITION UNTIL USED.

TAKE THREE QUART SIZE CANNING JARS AND PLACE THEM ASIDE, LABEL THEM 1, 2 AND 3.

BLEND SOME BONELESS CHEAP GREASY BEEF IN A BLENDER AND ADD AN EQUAL AMOUNT OF TAP WATER TO THE BEEF WHILE BLENDING.

POUR THE BEEF INTO EACH CANNING JAR, LEAVING ONE INCH FROM THE TOP.

ADD EACH SOIL SAMPLE, CAREFULLY NOTING IT'S INDIVIDUAL LOCATIONS AND LABEL THAT ON EACH JAR...

CAP EACH JAR AND SHAKE UNTIL WELL MIXED.

THE LIDS SHOULD BE PLACED ON FOR CANNING NOW, AND EACH JAR SHOULD BE PLACED IN A LARGE KETTLE FOR HEAT SHOCKING. LEAVE THE JARS IN THE BOILING WATER FOR 1 HOUR. LET THE GREASE RISE TO THE TOP TO SEAL THE AIR OUT. IF WHILE HEATING, THE JARS ARE NOT FULL ADD SOME MORE MEAT MUSH UNTIL THEY ARE FULL TO THE TOP.

THE CANNING RINGS SHOULD BE LOOSENED AS SOON AS THE JARS COOL DOWN, THE GERMS FORM A GAS THAT NEEDS TO BREATHE OUT OR THE JARS WILL EXPLODE WITH ROTTEN EGG ODORS. PLACE THESE JARS IN A WARM SHADED PLACE FOR 3-5 DAYS.

THE MEAT SHOULD HAVE TURNED BLACK WHERE THE GERMS ARE LOCATED. TEST THE POISON ON RATS OR CATS, ETC...MIX A LITTLE WITH SOME FOOD AND SHOVE IT DOWN THEIR THROAT AND IN A COUPLE DAYS THEY SHOULD BE DEAD, IF IT IS A GOOD BATCH.

IF IT IS A GOOD BATCH YOU CAN USE THE SPORES FROM THE BATCH THAT WORKED TO MAKE LOTS MORE ON A LARGER SCALE...JUST REMEMBER THAT AIR KILLS THE GERMS BUT NOT THE SPORES AND HEAT KILLS THE GERMS AND POISON BUT NOT THE SPORES. SO DO NOT HEAT THE POISON AFTER IT IS MADE....

BASIC SOLUTIONS ALSO KILL THE POISON.

HEAT KILLS THE POISON.

AIR KILLS THE GERMS.

HEAT KILLS THE GERMS.

HEAT DOES NOT KILL THE SPORES.

SLIGHTLY ACID SOLUTIONS DO NOT KILL THE GERMS, SPORES OR POISON.

IT IS BETTER TO MIX DMSO WITH A SLIGHTLY ACID SOLUTION FOR BETTER ASORBTION THROUGH THE SKIN.

50% PURIFICATION CAN BE ACHEIVED IF YOU ISOLATE THE BLACK MATERIAL MADE FROM THE BACTERIA AND PLACE IT IN A BEAKER

CONTAINING 3M HCL (10% CONCENTRATION)...THIS WILL CAUSE THE POISON TO PRECIPITATE TO THE BOTTOM OF THE BEAKER, THE PH MUST BE BETWEEN 3.5 AND 4. ALLOW THE BEAKER TO SIT OVERNIGHT AT ROOM TEMPERATURE. POUR OFF THE ACID AND PLACE THE LAST BIT CONTAINING THE POISON IN ANOTHER SMALLER JAR AND LET SIT AT 40 DEG F. NOW PLACE THIS IN A CENTRIFUGE AND SPIN AT 3500RPM FOR HALF AN HOUR...THE PELLETT IS VERY DANGEROUS BOTULIN POISON. MIX WITH LEMON JUICE AND DMSO AND THERE YOU HAVE IT...

INFO-LABS IS NOT RESPONSIBLE FOR USE OR MISUSE OF THIS INFORMATION. ANYONE ATTEMPTING TO MAKE THIS SUBSTANCE OR OTHERS LIKE IT ARE SOLELY LIABLE FOR ANY LEGALITIES PERTAINING.

DO NOT MAKE THIS MATERIAL, DO NOT USE THIS MATERIAL.
